

Successful Endovascular Stroke Rescue With Retrieval of an Embolized Calcium Fragment After Transcatheter Aortic Valve Replacement

Amir-Ali Fassa, MD; Mikael Mazighi, MD, PhD; Dominique Himbert, MD; Lydia Deschamps, MD; Gregory Ducrocq, MD; Adrian P. Cheong, MD; Jean-Pol Depoix, MD; Marie-Pierre Dilly, MD; Soleiman Alkholder, MD; Bruno Mourvillier, MD; Alec Vahanian, MD

A 90-year-old symptomatic woman with a critical aortic stenosis was referred for transcatheter aortic valve replacement (TAVR). The procedure was performed under locoregional anesthesia from a right femoral approach, with the successful implantation of a 23-mm CoreValve (Medtronic Inc, Minneapolis, MN). Echocardiographic assessment after TAVR showed a mean transprosthetic gradient of 16 mmHg and trace paravalvular regurgitation. After percutaneous closure of the right femoral artery, the patient suddenly became unresponsive. After prompt intubation, emergency cerebral MRI was performed to assess the presence of reversible ischemia and exclude parenchymal hemorrhage, showing partial occlusion of the right middle cerebral artery with ischemia in the corresponding territory (Figure 1). Conventional cerebral angiography confirmed partial M1-M2 occlusion of the right middle cerebral artery (Figure 2A; Movie I in the Data Supplement). Complete revascularization was achieved using a 4.0×20 mm Solitaire FR retrievable stent (ev3, Irvine, CA) with capture of the embolic material (Figure 2B; Movies II and III in the Data Supplement). The onset-to-reperfusion delay was 150 minutes.

Histopathologic assessment of the recovered debris (2×6 mm) revealed a calcium fragment that most likely detached from the native aortic valve or the aortic wall (Figure 3).

The patient was extubated on the following day. She was fully conscious and orientated, with normal neurological status (modified Rankin Scale score, 0). She had an uneventful recovery and was discharged 9 days after the procedure.

Discussion

Stroke occurs during or <24 hours of TAVR in ≈3% of cases and is associated with a 12-fold increase in 30-day mortality.¹ This complication is presumably related to manipulation of large-bore catheters across the aorta and the calcified aortic valve. A recent histopathologic study on embolized debris

captured during TAVR with a filter-based cerebral protection device revealed either thrombotic material or embolized tissue debris derived most likely from the native aortic valve leaflets or aortic wall.²

Endovascular intervention for stroke due to catheter-related thrombus during TAVR has been previously reported.³ Although mechanical retrieval of nonthrombotic material from cerebral arteries after acute ischemic stroke has also been described,⁴ the present case is the first report of retrieval of embolized tissue material presumably originating from the native aortic valve or aortic wall after TAVR. Furthermore, the Solitaire FR retrievable stent has been proven effective in the treatment of acute ischemic stroke, with high rates of successful recanalization and favorable neurological outcomes during follow-up, as well as a low incidence of procedure-related complications (such as distal emboli to a new territory or vessel dissection or perforation).⁵ In the setting of stroke complicating TAVR, mechanical endovascular revascularization with retrievable stents may be the treatment of choice, because sole intravenous or intra-arterial thrombolysis is contraindicated in fully heparinized patients and may be ineffective in the presence of embolized tissue material.

Ultimately, the present case underscores the advantage of locoregional anesthesia during TAVR for immediate diagnosis of neurological events, because interventional rescue could be potentially delayed when patients are intubated and sedated.

Disclosures

Dr Himbert is a proctor for Medtronic Inc. Dr Vahanian received speaker's fees from Medtronic Inc. The other authors have no conflicts to report.

References

1. Stortecky S, Windecker S, Pilgrim T, Heg D, Buellesfeld L, Khattab AA, Huber C, Gloekler S, Nietlisbach F, Mattle H, Jüni P, Wenaweser

Received October 18, 2013; accepted December 5, 2013.

From the Departments of Cardiology (A.-A.F., D.H., G.D., A.P.C., A.V.), Neurology and Stroke Centre (M.M.), Pathology (L.D.), Anesthesiology (J.-P.D., M.-P.D.), Cardiac Surgery (S.A.), and Intensive Care and Infectious Diseases (B.M.), Bichat-Claude-Bernard Hospital, Assistance-Publique-Hôpitaux de Paris, Université Paris VII, Paris, France.

The Data Supplement is available at <http://circinterventions.ahajournals.org/lookup/suppl/doi:10.1161/CIRCINTERVENTIONS.113.000995/-/DC1>.

Correspondence to Amir-Ali Fassa, MD, Department of Cardiology, Bichat Hospital, 46 rue Henri-Huchard, 75018 Paris, France. E-mail amir.fassa@gmail.com

(*Circ Cardiovasc Interv.* 2014;7:125-126.)

© 2014 American Heart Association, Inc.

Circ Cardiovasc Interv is available at <http://circinterventions.ahajournals.org>

DOI: 10.1161/CIRCINTERVENTIONS.113.000995

P. Cerebrovascular accidents complicating transcatheter aortic valve implantation: frequency, timing and impact on outcomes. *EuroIntervention*. 2012;8:62–70.

2. Van Mieghem NM, Schipper ME, Ladich E, Faqiri E, van der Boon R, Randjgari A, Schultz C, Moelker A, van Geuns RJ, Otsuka F, Serruys PW, Virmani R, de Jaegere PP. Histopathology of embolic debris captured during transcatheter aortic valve replacement. *Circulation*. 2013;127:2194–2201.

3. Salinas P, Moreno R, Frutos R, Lopez-Sendon JL. Neurovascular rescue for thrombus-related embolic stroke during transcatheter aortic valve implantation. *J Am Coll Cardiol Cardiovasc Interv*. 2013;6:981–982.

4. Marder VJ, Chute DJ, Starkman S, Abolian AM, Kidwell C, Liebeskind D, Ovbiagele B, Vinuela F, Duckwiler G, Jahan R, Vespa PM, Selco S, Rajajee V, Kim D, Sanossian N, Saver JL. Analysis of thrombi retrieved from cerebral arteries of patients with acute ischemic stroke. *Stroke*. 2006;37:2086–2093.

5. Saver JL, Jahan R, Levy EI, Jovin TG, Baxter B, Nogueira RG, Clark W, Budzik R, Zaidat OO; SWIFT Trialists. Solitaire flow restoration device versus the Merci Retriever in patients with acute ischaemic stroke (SWIFT): a randomised, parallel-group, non-inferiority trial. *Lancet*. 2012;380:1241–1249.

KEY WORD: stroke




Figure 1. **A**, Diffusion-weighted MRI showing ischemia (faint hypersignal; arrow) in the territory of the right middle cerebral artery. **B** to **D**, Magnetic resonance angiography showing partial occlusion of the right middle cerebral artery (arrowhead) on 3-dimensional time-of-flight sequences. **C**, Close-up of the boxed area in **B**.




Figure 2. **A**, Conventional cerebral angiography confirming partial M1-M2 occlusion of the right middle cerebral artery (arrowhead). **B**, Final angiography after retrieval of the embolized debris showing complete patency of the right middle cerebral artery.




Figure 3. **A**, Debris recovered during endovascular revascularization. The distance between the lines of the scale on the left is 1 mm. **B**, Histopathologic assessment (hematoxylin and eosin stained) showing pure calcific material. **C** and **D**, High-power magnification of the boxed areas in **B**.

Successful Endovascular Stroke Rescue With Retrieval of an Embolized Calcium Fragment After Transcatheter Aortic Valve Replacement

Amir-Ali Fassa, Mikael Mazighi, Dominique Himbert, Lydia Deschamps, Gregory Ducrocq, Adrian P. Cheong, Jean-Pol Depoix, Marie-Pierre Dilly, Soleiman Alkhoder, Bruno Mourvillier and Alec Vahanian

Circ Cardiovasc Interv. 2014;7:125-126

doi: 10.1161/CIRCINTERVENTIONS.113.000995

Circulation: Cardiovascular Interventions is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2014 American Heart Association, Inc. All rights reserved.

Print ISSN: 1941-7640. Online ISSN: 1941-7632

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circinterventions.ahajournals.org/content/7/1/125>

Data Supplement (unedited) at:

<http://circinterventions.ahajournals.org/content/suppl/2014/02/12/CIRCINTERVENTIONS.113.000995.DC1>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Interventions* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation: Cardiovascular Interventions* is online at:
<http://circinterventions.ahajournals.org/subscriptions/>

SUPPLEMENTAL MATERIAL

Movie I. Conventional cerebral angiography showing partial M1-M2 occlusion of the right middle cerebral artery.

Movie II. Deployment of the Solitaire FR retrievable stent (ev3, Irvine, CA), with restoration of flow (the three dots indicate the distal marker of the device).

Movie III. Final angiography after retrieval of the embolized debris showing complete patency of the right middle cerebral artery.